




School Nursing Life-Threatening Allergy IHP

() Epinephrine 0.15mg () Antihistamine- name/do () If checked, OK for stu	treat a severe reaction. Use Epinephrine. () Epinephrine 0.3mg ose udent to carry/self-administer. Relationship:	
() Epinephrine 0.15mg () Antihistamine- name/do () If checked, OK for stu	treat a severe reaction. Use Epinephrine. () Epinephrine 0.3mg ose udent to carry/self-administer. Relationship:	_ Date:
lepend on antihistamines to a () Epinephrine 0.15mg () Antihistamine- name/do () If checked, OK for stu emergency contact:	treat a severe reaction. Use Epinephrine. () Epinephrine 0.3mg ose udent to carry/self-administer.	
() Epinephrine 0.15mg () Antihistamine- name/do () If checked, OK for stu	treat a severe reaction. Use Epinephrine. () Epinephrine 0.3mg ose udent to carry/self-administer.	Situation. Do not
() Epinephrine 0.15mg () Antihistamine- name/do () If checked, OK for stu	treat a severe reaction. Use Epinephrine. () Epinephrine 0.3mg ose udent to carry/self-administer.	Situation. Do not
() Epinephrine 0.15mg () Antihistamine- name/do () If checked, OK for stu	treat a severe reaction. Use Epinephrine. () Epinephrine 0.3mg ose udent to carry/self-administer.	Situation. Do not
() Epinephrine 0.15mg () Antihistamine- name/do	treat a severe reaction. Use Epinephrine. () Epinephrine 0.3mg ose	Situation. Do not
lepend on antihistamines to to (() Epinephrine 0.15mg	treat a severe reaction. Use Epinephrine. () Epinephrine 0.3mg	Situation. Do not
		Situation. Do not
MOUTH: THROAT*: SKIN: GUT: LUNG*: HEART*:	itching, swelling of lips and/or tongue itching, tightness/closure, hoarseness, trouble breathir itching, hives, redness, swelling vomiting, diarrhea, cramps Shortness of breath, cough, wheeze weak pulse, dizziness, passing out, pale, faint	
ive epinephrine immediately ive epinephrine immediately	for ANY symptoms if the allergen was likely eaten if the allergen was definitely eaten, even if no sympton	ns are noted.
the school with a supply of medica the school with the written doctor' edication will not be administered ne school of any medical changes the school with this signed conser	i's instructions for medication administration during school hours. until signed doctors instructions are at school s. ent form annually and when changes in medication occur.	acy.
ing administrator or his/her design onsent for the exchange of information hare, if necessary, this information	nee administer the medication or procedure listed below as nation between the school and my child's health care provider. on with school personnel who may be involved with the welfare of	
D BY PARENT/GUARDIAN	ems are to be brought to the school by the parent/guardian.	
	of	In of

Fennville Public Schools
Hamilton Public Schools
Holland Christian Schools
Holland Public Schools
West Ottawa Public Schools



School Nursing Life-Threatening Allergy IHP

EPIPEN® (EPINEPHRINE) AUTO-INJECTOR DIRECTIONS

- 1. Remove the EpiPen Auto-Injector from the plastic carrying case.
- 2. Pull off the blue safety release cap.
- 3. Swing and firmly push orange tip against mid-outer thigh.
- 4. Hold for approximately 10 seconds.
- 5. Remove and massage the area for 10 seconds.

